

KIC 002983000

Q1-17 DR25 TCE Parameters

TCE	Run Type	KOI?	Period (Days)	Epoch (BKJD)	Depth (ppm)	Duration (Hours)	MES	SNR	R_{\star} (R_{\odot})	T_{\star} (K)	R_p (R_{\oplus})	S_p (S_{\oplus})
002983000-01	OBS	No	602.327552	156.477752	1132.0	17.608	19.2	17.9	0.97	6108	3.33	0.62

Robovetter Results

TCE	Run Type	Disp	Score	N	S	C	E	Comments
002983000-01	OBS	FP	0.00	1	0	1	0	INDIV_TRANS_CHASES_MARSHALL—LPP_DV—INCONSISTENT_TRANS—CENT_FEW_DIFFS—HALO_GHOST

Notes: OBS = Observed. INJ = Injected. INV = Inverted. SCR = Scrambled.

N = Not Transit-Like. S = Stellar Eclipse. C = Centroid Offset. E = Ephemeris Match.


See http://exoplanetarchive.ipac.caltech.edu/docs/API_kepcandidate_columns.html#proj_disp_col for comment definitions.

Ephemeris Match Information For 002983000-01

No Significant Match Found

DV One-Page Summary


KIC: 2983000 Candidate: 1 of 1 Period: 602.328 d




Software Revision: svn+ssh://murzim/repo/soc/tags/release/9.3.42@60958 -- Date Generated: 30-Jan-2016 15:04:53 Z

This Data Validation Report Summary was produced in the Kepler Science Operations Center Pipeline at NASA Ames Research Center

TCE 002983000-01, PDC Light Curves




TCE 002983000-01




DV Odd/Even

TCE 002983000-01




ALT Odd/Even

TCE 002983000-01




Non-Whitened Vs. Whitened Light Curve

Planet 1 : Phased Unwhitened Flux Time Series (Fit Epoch/Period)




Planet 1 : Phased Whitened Flux Time Series (Fit Epoch/Period)




PDC Quarter-Phased Transit Curves

TCE 002983000-01 P=602.327552 Days $T_0=156.477752$ (BKJD)




DV Quarter-Phased Transit Curves

TCE 002983000-01 P=602.327552 Days $T_0=156.477752$ (BKJD)



Alt. Detrend Quarter-Phased Transit Curves


TCE 002983000-01 P=602.265559 Days $T_0=156.537717$ (BKJD)



DV Model-Shift Uniqueness Test

002983000-01, P = 602.327552 Days, E = 156.477752 Days


Pri	Sec	Ter	Pos	FA ₁	FA ₂	F _{Red}	Pri-Ter	Pri-Pos	Sec-Ter	Sec-Pos	Odd-Evn	DMM	Shape	TAT
12.7	6.12	5.57	5.55	5.41	3.23	1.35	7.11	7.13	0.54	0.57	28.2	4.66	0.30	4.75



Alt Model-Shift Uniqueness Test

002983000-01, P = 602.265559 Days, E = 156.537717 Days

Pri	Sec	Ter	Pos	FA ₁	FA ₂	F _{Red}	Pri-Ter	Pri-Pos	Sec-Ter	Sec-Pos	Odd-Evn	DMM	Shape	TAT
15.0	5.75	5.08	4.84	5.41	3.23	1.36	9.91	10.1	0.67	0.91	24.8	3.20	0.24	0.68



Stellar Parameters For KIC 002983000

	$T_{\text{eff}}(K)$	$\log(g)$	[Fe/H]	$R (R_{\odot})$	$M(M_{\odot})$	$p_{\star} (\text{g}\cdot\text{cm}^{-3})$
	6108^{+183}_{-201}	$4.443^{+0.084}_{-0.196}$	$-0.340^{+0.300}_{-0.300}$	$0.974^{+0.280}_{-0.120}$	$0.959^{+0.128}_{-0.102}$	$1.463^{+0.525}_{-0.743}$
	+3%/-3%	+2%/-4%	+88%/-88%	+29%/-12%	+13%/-11%	+36%/-51%
Source	PHO1	KIC0	KIC0	DSEP		

KIC = Kepler Input Catalog; PHO = Photometry; SPE = Spectroscopy; AST = Asteroseismology
 TRA = Transits; DESP = Dartmouth Models; MULT = Multiple Models

Secondary Eclipse Parameters for KIC 002983000-01 / KOI

Detrend	Depth (ppm)	$R_p (R_{\oplus})$	$T_{\text{max}} (K)$	$T_{\text{obs}} (K)$	A_{obs}
DV	-216 ± 35	$3.46^{+0.71}_{-0.65}$	321^{+23}_{-16}	4397^{+349}_{-287}	19044^{+9607}_{-6759}
Alt.	-210 ± 37	$3.10^{+0.68}_{-0.57}$	321^{+22}_{-16}	4537^{+420}_{-332}	22627^{+12465}_{-8092}

T_{max} = Theoretical Maximum Planetary Temperature

T_{obs} = Observed Planetary Temperature (Assuming $A=0.3$)

A_{obs} = Observed Albedo (Assuming $T=0$)

If a secondary eclipse is present, the system is likely an EB if $T_{\text{obs}} \gg T_{\text{max}}$ AND $A_{\text{obs}} \gg 1.0$


DV Centroid Data

Supplemental centroid analysis for 002983000-01. Kepler magnitude: 14.63. Transit SNR 17.88

There are 1 quarters with good PRF difference image offsets


The direct PRF centroid is offset from the target star catalog position by about 0.38 arcsec

	Distance in arcsec	Distance / σ	Δ RA	Δ Dec
PRF-fit source offset from OOT	2.146 ± 1.836	1.17	-0.483 ± 0.171	2.091 ± 1.884
PRF-fit source offset from KIC position	2.284 ± 1.814	1.26	-0.539 ± 0.202	2.220 ± 1.866
photometric centroid source offset	0.21 ± 0.37	0.57	-0.21 ± 0.37	0.02 ± 0.45



Centroid source offsets from the target star reconstructed from PRF and photometric centroids. Sky blue crosses: good quarterly centroid offsets; Vermillion crosses: bad quarterly centroid offsets; magenta cross: average over quarters. Length of the crosses: one- σ uncertainty. Blue circle: three- σ . Red *: target star. Blue *: Other stars. Text next to a star gives its KIC ID and kepmag. KIC IDs > 15,000,000 are from the UKIRT catalog.

white ×: KIC target position; +: OOT centroid; △: difference centroid. red ✕: large negative pixel value.



white \times : KIC target position; +: OOT centroid; \triangle : difference centroid. red \times : large negative pixel value.

Q5 no difference image



Q5 no OOT image



Q6 no difference image



Q6 no OOT image




Q7 no difference image




Q7 no OOT image



Q8 difference image



Q8 OOT image



white \times : KIC target position; $+$: OOT centroid; \triangle : difference centroid. red \times : large negative pixel value.




white \times : KIC target position; +: OOT centroid; \triangle : difference centroid. red \times : large negative pixel value.


Q13 no difference image




Q13 no OOT image



Q14 difference image. Poor Quality




Q14 OOT image



Q15 no difference image




Q15 no OOT image




Q16 no difference image



Q16 no OOT image



white \times : KIC target position; $+$: OOT centroid; \triangle : difference centroid. red \times : large negative pixel value.



UKIRT Image

Declination

