

KIC 005215576

Q1-17 DR25 TCE Parameters

TCE	Run Type	KOI?	Period (Days)	Epoch (BKJD)	Depth (ppm)	Duration (Hours)	MES	SNR	R_{\star} (R_{\odot})	T_{\star} (K)	R_p (R_{\oplus})	S_p (S_{\oplus})
005215576-01	OBS	No	40.588515	137.443465	171.7	62.539	7.9	12.1	0.80	5667	2.13	13.94

Robovetter Results

TCE	Run Type	Disp	Score	N	S	C	E	Comments
005215576-01	OBS	FP	0.00	1	0	0	0	LPP_DV—LPP_ALT—CENT_FEW_DIFFS

Notes: OBS = Observed. INJ = Injected. INV = Inverted. SCR = Scrambled.

N = Not Transit-Like. S = Stellar Eclipse. C = Centroid Offset. E = Ephemeris Match.


See http://exoplanetarchive.ipac.caltech.edu/docs/API_kepcandidate_columns.html#proj_disp_col for comment definitions.

Ephemeris Match Information For 005215576-01

No Significant Match Found

DV One-Page Summary


KIC: 5215576 Candidate: 1 of 1 Period: 40.589 d




Software Revision: svn+ssh://murzim/repo/soc/tags/release/9.3.42@60958 -- Date Generated: 29-Jan-2016 08:50:06 Z

This Data Validation Report Summary was produced in the Kepler Science Operations Center Pipeline at NASA Ames Research Center

TCE 005215576-01, PDC Light Curves




TCE 005215576-01




DV Odd/Even

TCE 005215576-01




ALT Odd/Even

TCE 005215576-01




Non-Whitened Vs. Whitened Light Curve

Planet 1 : Phased Unwhitened Flux Time Series (Fit Epoch/Period)




Planet 1 : Phased Whitened Flux Time Series (Fit Epoch/Period)




PDC Quarter-Phased Transit Curves

TCE 005215576-01 P= 40.588515 Days $T_0=137.443465$ (BKJD)




DV Quarter-Phased Transit Curves

TCE 005215576-01 P= 40.588515 Days $T_0=137.443465$ (BKJD)



Alt. Detrend Quarter-Phased Transit Curves


TCE 005215576-01 P= 40.595497 Days $T_0=137.234128$ (BKJD)



DV Model-Shift Uniqueness Test

005215576-01, P = 40.588515 Days, E = 96.854950 Days


Pri	Sec	Ter	Pos	FA ₁	FA ₂	F _{Red}	Pri-Ter	Pri-Pos	Sec-Ter	Sec-Pos	Odd-Evn	DMM	Shape	TAT
30.4	11.7	4.83	8.58	4.66	1.85	3.32	25.6	21.8	6.87	3.11	3.84	1.34	0.41	0.86



Alt Model-Shift Uniqueness Test

005215576-01, P = 40.595497 Days, E = 96.638631 Days

Pri	Sec	Ter	Pos	FA ₁	FA ₂	F _{Red}	Pri-Ter	Pri-Pos	Sec-Ter	Sec-Pos	Odd-Evn	DMM	Shape	TAT
11.5	2.51	2.19	1.94	4.63	1.80	1.14	9.33	9.58	0.31	0.56	2.45	1.41	0.31	1.30



Stellar Parameters For KIC 005215576

	$T_{\text{eff}}(K)$	$\log(g)$	[Fe/H]	$R (R_{\odot})$	$M(M_{\odot})$	$p_{\star} (\text{g}\cdot\text{cm}^{-3})$
	5667^{+183}_{-167}	$4.483^{+0.130}_{-0.159}$	$-0.740^{+0.300}_{-0.300}$	$0.803^{+0.165}_{-0.110}$	$0.714^{+0.102}_{-0.036}$	$1.944^{+1.086}_{-0.812}$
	+3%/-3%	+3%/-4%	+41%/-41%	+21%/-14%	+14%/-5%	+56%/-42%
Source	PHO1	KIC0	KIC0	DSEP		

KIC = Kepler Input Catalog; PHO = Photometry; SPE = Spectroscopy; AST = Asteroseismology
 TRA = Transits; DESP = Dartmouth Models; MULT = Multiple Models

Secondary Eclipse Parameters for KIC 005215576-01 / KOI

Detrend	Depth (ppm)	$R_p (R_{\oplus})$	$T_{\text{max}} (K)$	$T_{\text{obs}} (K)$	A_{obs}
DV	-64 ± 6	$4.38^{+4.01}_{-3.13}$	689^{+43}_{-38}	3005^{+1410}_{-502}	82^{+973}_{-60}
Alt.	-13 ± 5	$3.27^{+3.51}_{-2.34}$	693^{+44}_{-40}	2542^{+1130}_{-412}	27^{+311}_{-21}

T_{max} = Theoretical Maximum Planetary Temperature

T_{obs} = Observed Planetary Temperature (Assuming $A=0.3$)

A_{obs} = Observed Albedo (Assuming $T=0$)

If a secondary eclipse is present, the system is likely an EB if $T_{\text{obs}} \gg T_{\text{max}}$ AND $A_{\text{obs}} \gg 1.0$


DV Centroid Data

Supplemental centroid analysis for 005215576-01. Kepler magnitude: 14.64. Transit SNR 12.14

There are 2 quarters with good PRF difference image offsets





The direct PRF centroid is offset from the target star catalog position by about 0.09 arcsec

	Distance in arcsec	Distance / σ	Δ RA	Δ Dec
PRF-fit source offset from OOT	0.893 ± 1.358	0.66	0.864 ± 1.383	0.225 ± 0.917
PRF-fit source offset from KIC position	0.899 ± 1.342	0.67	0.884 ± 1.354	0.163 ± 0.936
photometric centroid source offset	0.27 ± 0.85	0.32	0.04 ± 0.95	-0.27 ± 0.85







Centroid source offsets from the target star reconstructed from PRF and photometric centroids. Sky blue crosses: good quarterly centroid offsets; Vermillion crosses: bad quarterly centroid offsets; magenta cross: average over quarters. Length of the crosses: one- σ uncertainty. Blue circle: three- σ . Red *: target star. Blue *: Other stars. Text next to a star gives its KIC ID and kepmag. KIC IDs > 15,000,000 are from the UKIRT catalog.



white \times : KIC target position; +: OOT centroid; \triangle : difference centroid. red \times : large negative pixel value.



white \times : KIC target position; +: OOT centroid; \triangle : difference centroid. red \times : large negative pixel value.



white \times : KIC target position; +: OOT centroid; \triangle : difference centroid. red \times : large negative pixel value.



white ×: KIC target position; +: OOT centroid; △: difference centroid. red ✕: large negative pixel value.


Q13 no difference image




Q13 no OOT image



Q14 difference image. Poor Quality



Q14 OOT image



Q15 no difference image



Q15 no OOT image




Q16 no difference image



Q16 no OOT image



white \times : KIC target position; $+$: OOT centroid; \triangle : difference centroid. red \times : large negative pixel value.



UKIRT Image

Declination

