

KIC 006676298

Q1-17 DR25 TCE Parameters

TCE	Run Type	KOI?	Period (Days)	Epoch (BKJD)	Depth (ppm)	Duration (Hours)	MES	SNR	R_{\star} (R_{\odot})	T_{\star} (K)	R_p (R_{\oplus})	S_p (S_{\oplus})
006676298-01	OBS	No	555.524490	450.288695	516.3	3.983	7.3	7.7	0.95	6075	2.31	0.62

Robovetter Results

TCE	Run Type	Disp	Score	N	S	C	E	Comments
006676298-01	OBS	FP	0.00	1	0	0	0	INDIV_TRANS_CHASES_MARSHALL—LPP_DV—LPP_ALT—ALL_TRANS_CHASES—MOD_NONUNIQ_ALT—MOD_TER_ALT—MOD_POS_ALT— INCONSISTENT_TRANS—CENT_FEW_MEAS

Notes: OBS = Observed. INJ = Injected. INV = Inverted. SCR = Scrambled.

N = Not Transit-Like. S = Stellar Eclipse. C = Centroid Offset. E = Ephemeris Match.


See http://exoplanetarchive.ipac.caltech.edu/docs/API_kepcandidate_columns.html#proj_disp_col for comment definitions.

Ephemeris Match Information For 006676298-01

No Significant Match Found

DV One-Page Summary


KIC: 6676298 Candidate: 1 of 1 Period: 555.524 d




Software Revision: svn+ssh://murzim/repo/soc/tags/release/9.3.42@60958 -- Date Generated: 31-Jan-2016 14:58:42 Z

This Data Validation Report Summary was produced in the Kepler Science Operations Center Pipeline at NASA Ames Research Center

TCE 006676298-01, PDC Light Curves




TCE 006676298-01




DV Odd/Even

TCE 006676298-01




ALT Odd/Even

TCE 006676298-01




Non-Whitened Vs. Whitened Light Curve

Planet 1 : Phased Unwhitened Flux Time Series (Fit Epoch/Period)




Planet 1 : Phased Whitened Flux Time Series (Fit Epoch/Period)




PDC Quarter-Phased Transit Curves

TCE 006676298-01 $P=555.524490$ Days $T_0=450.288695$ (BKJD)




DV Quarter-Phased Transit Curves

TCE 006676298-01 P=555.524490 Days $T_0=450.288695$ (BKJD)



Alt. Detrend Quarter-Phased Transit Curves


TCE 006676298-01 P=555.520384 Days $T_0=450.286826$ (BKJD)



DV Model-Shift Uniqueness Test

006676298-01, P = 555.524490 Days, E = 450.288695 Days


Pri	Sec	Ter	Pos	FA ₁	FA ₂	F _{Red}	Pri-Ter	Pri-Pos	Sec-Ter	Sec-Pos	Odd-Evn	DMM	Shape	TAT
7.31	4.64	4.02	4.56	5.65	3.60	1.07	3.29	2.75	0.62	0.07	1.83	0.91	0.38	1.55



Alt Model-Shift Uniqueness Test

006676298-01, P = 555.520384 Days, E = 450.286826 Days

Pri	Sec	Ter	Pos	FA ₁	FA ₂	F _{Red}	Pri-Ter	Pri-Pos	Sec-Ter	Sec-Pos	Odd-Evn	DMM	Shape	TAT
6.18	4.77	4.32	4.72	5.66	3.61	0.96	1.86	1.46	0.45	0.05	1.83	0.90	0.43	0.91



Stellar Parameters For KIC 006676298

	$T_{\text{eff}}(K)$	$\log(g)$	[Fe/H]	R (R_{\odot})	M (M_{\odot})	p_{\star} ($\text{g}\cdot\text{cm}^{-3}$)
	6075^{+169}_{-190}	$4.495^{+0.050}_{-0.212}$	$-0.160^{+0.300}_{-0.300}$	$0.952^{+0.291}_{-0.097}$	$1.033^{+0.140}_{-0.140}$	$1.689^{+0.444}_{-0.847}$
	+3%/-3%	+1%/-5%	+188%/-188%	+31%/-10%	+14%/-14%	+26%/-50%
Source	PHO1	KIC0	KIC0	DSEP		

KIC = Kepler Input Catalog; PHO = Photometry; SPE = Spectroscopy; AST = Asteroseismology
 TRA = Transits; DESP = Dartmouth Models; MULT = Multiple Models

Secondary Eclipse Parameters for KIC 006676298-01 / KOI

Detrend	Depth (ppm)	R_p (R_{\oplus})	T_{max} (K)	T_{obs} (K)	A_{obs}
DV	-258 ± 56	$3.11^{+2.51}_{-2.08}$	323^{+23}_{-15}	4660^{+3370}_{-887}	$25403^{+199840}_{-17780}$
Alt.	-262 ± 55	$2.78^{+2.32}_{-1.66}$	324^{+24}_{-15}	4895^{+3175}_{-981}	$30918^{+183120}_{-21516}$

T_{max} = Theoretical Maximum Planetary Temperature

T_{obs} = Observed Planetary Temperature (Assuming $A=0.3$)

A_{obs} = Observed Albedo (Assuming $T=0$)

If a secondary eclipse is present, the system is likely an EB if $T_{obs} \gg T_{max}$ AND $A_{obs} \gg 1.0$


DV Centroid Data

Supplemental centroid analysis for 006676298-01. Kepler magnitude: 14.76. Transit SNR 7.65

There are 1 quarters with good PRF difference image offsets

The direct PRF centroid is offset from the target star catalog position by about 0.25 arcsec

	Distance in arcsec	Distance / σ	Δ RA	Δ Dec
PRF-fit source offset from OOT	2.901 ± 0.321	9.03	1.966 ± 0.334	2.134 ± 0.310
PRF-fit source offset from KIC position	2.711 ± 0.323	8.40	1.950 ± 0.334	1.883 ± 0.310
photometric centroid source offset	4.27 ± 2.66	1.61	-0.48 ± 1.80	4.24 ± 2.67






Centroid source offsets from the target star reconstructed from PRF and photometric centroids. Sky blue crosses: good quarterly centroid offsets; Vermillion crosses: bad quarterly centroid offsets; magenta cross: average over quarters. Length of the crosses: one- σ uncertainty. Blue circle: three- σ . Red *: target star. Blue *: Other stars. Text next to a star gives its KIC ID and kepmag. KIC IDs > 15,000,000 are from the UKIRT catalog.

white \times : KIC target position; $+$: OOT centroid; \triangle : difference centroid. red \times : large negative pixel value.



white ×: KIC target position; +: OOT centroid; △: difference centroid. red ✕: large negative pixel value.



white \times : KIC target position; $+$: OOT centroid; \triangle : difference centroid. red \times : large negative pixel value.

Q9 no difference image



Q9 no OOT image




Q10 no difference image




Q10 no OOT image



Q11 difference image. Poor Quality



Q11 OOT image



Q12 no difference image




Q12 no OOT image




white \times : KIC target position; $+$: OOT centroid; \triangle : difference centroid. red \times : large negative pixel value.



white \times : KIC target position; $+$: OOT centroid; \triangle : difference centroid. red \times : large negative pixel value.



fluxWeightedCentroids, Planet 1 of 1



UKIRT Image

Declination

